

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/14, 9/16, 9/22, 9/24, 9/30, 9/36, 9/42, A01N 37/00, 37/12		A1	(11) International Publication Number: WO 00/12064
			(43) International Publication Date: 9 March 2000 (09.03.00)
(21) International Application Number: PCT/US99/19847			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 27 August 1999 (27.08.99)			
(30) Priority Data: 09/143,167 28 August 1998 (28.08.98) US			
(71) Applicant (for all designated States except US): ANDRX PHARMACEUTICALS, INC. [US/US]; Suite 201, 4001 S.W. 47th Avenue, Fort Lauderdale, FL 33314 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): CHEN, Chih-Ming [US/US]; 10680 S.W. 40th Manor, Davie, FL 33328 (US). CHOU, Joseph [US/US]; 5755 N.W. 54th Place, Coral Springs, FL 33067 (US). KOSITPRAPA, Unchalee [TH/US]; Apartment 206C, 4163 S.W. 67th Avenue, Fort Lauderdale, FL 33314 (US).			
(74) Agent: COSTIGAN, James, V.; Hedman, Gibson & Costigan, P.C., 1185 Avenue of the Americas, New York, NY 10036 (US).			Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: OMEPRAZOLE FORMULATION			
(57) Abstract <p>A stable pharmaceutical composition of omeprazole for oral administration which consists essentially of: (a) a core of Omeprazole or a pharmaceutically equivalent salt, a filler and an alkaline material selected from the group consisting of lysine and arginine; and (b) a single layer of coating on said core which comprises a layer of an enteric coating agent which is applied from an organic solvent based system.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

OMEPRAZOLE FORMULATIONBACKGROUND OF THE INVENTION

The present invention relates to a stable formulation of omeprazole. It is well known that omeprazole is sensitive to acidic conditions and after contact with an acid, omeprazole will degrade and will not function in its intended manner. Initially, alkaline materials were added to a core of omeprazole and later an enteric coating was applied over the core to prevent the omeprazole from contacting the acidic pH conditions of the stomach. This approach is satisfactory if the product is administered within a short time after it is manufactured but if the product is stored under ambient conditions, the acidic residue of the enteric coating appears to degrade the omeprazole before it is administered to a patient. To solve this problem, the prior art has used a separate layer of a coating agent to coat a pellet core which contains omeprazole and an alkaline material which is thereafter coated with the enteric coating. This technique is described in U.S. 4,786,505. In addition WO 96/24338 discloses the use of an in situ formed interlayer that is based on the reaction of an aqueous enteric coating material with an alkaline material in the core.

This dual layer coating technique requires the application of two separate functional coating operations which increases the length of the manufacturing process and the cost of the product. The applicants have surprisingly discovered a coating system which avoids the need to use a coating layer to separate the omeprazole core from the enteric coating layer in an omeprazole dosage form. The separate coating system is based on the combined use of an enteric coating agent which is applied to a pelletized core or a granular core of omeprazole as a suspension

in a suitable solvent.

The applicants have also surprisingly discovered that arginine or lysine can be used as a pH stabilizing agent.

5

SUMMARY OF THE INVENTION

The present invention provides a novel stable pharmaceutical composition of omeprazole for oral
10 administration which consists essentially of:

- (a) a core of omeprazole or a pharmaceutically equivalent salt, a filler and an alkaline material selected from the group consisting of lysine and arginine; and
- 15 (b) a single layer of coating on said core which comprises a layer of an enteric coating agent applied from an organic based solvent coating system.

The core of the pharmaceutical composition can be in the form of a compressed tablet which is further
20 comprised essentially of a surface active agent, and a binder. Alternatively, the pharmaceutical composition can have a pelleted core which is further comprised essentially of an inert core component, a surface active agent and a binder.

25 Accordingly, it is a primary object of this invention to provide a pharmaceutical dosage formulation of omeprazole which is stable upon prolonged storage, is stable when administered to a patient and is capable of providing the desired
30 therapeutic effect.

It is also an object of this invention to provide a pharmaceutical dosage form of omeprazole which is bioequivalent to dosage forms of omeprazole which have an intermediate layer of an inert coating
35 material.

It is also an object of this invention to provide

a stable dosage form of omeprazole which may be produced without the need to provide an intermediate coating layer that separates the omeprazole containing core from the enteric coating layer.

5 These and other objects of the invention will become apparent from a review of the appended specification.

DETAILED DESCRIPTION OF THE INVENTION

10

The omeprazole formulation of the invention is preferably based on a core of omeprazole or pharmaceutically equivalent salt, a filler and an alkaline material selected from the group consisting of arginine or lysine; and a single layer of coating on said core which comprises a layer of an enteric coating agent applied from an organic solvent based system. The Omeprazole core can either be pelleted or tabletted as described herein.

20

In the case of both the pelleted form and the tabletted form of the core a filler is used. A filler is used as a granulation substrate. Sugars such as lactose, dextrose, sucrose, maltose, or microcrystalline cellulose and the like may be used as fillers in either the pellet or the granulation composition. In the case of the pelleted form the filler may comprise from 20 to 90wt% and preferably 65-85wt% based on the total weight of the drug layer composition. In the case of the tabletted form the filler may comprise from 20 to 60wt% and preferably 20 to 40wt% based on the total weight of the granulation. In the case of the tabletted form of the invention a tablet disintegrant may be added which comprises corn starch, potato starch, croscarmellose sodium, crospovidone and sodium starch glycolate in an effective amount. An effective amount which may be

25

30

35

from 3 to 10wt% based on the total weight of the granulation.

In the case of both the tabletted form and the pelleted form of the core an alkaline agent that is either lysine or arginine is used as a stabilizer. In the case of the tabletted form a level of from 20 to 60wt% and preferably 30 to 55wt% based on the weight of the granulation may be employed. In the case of the pelleted form a level of from 0.5 to 10wt% and preferably 1 to 3wt% based on the weight of the pellet may be employed.

In the case of both the pelleted form and the tabletted form of the invention an enteric coating agent is placed over the core. In both cases the enteric coating may comprise an acid resisting material which resists acid up to a pH of above about 5.0 or higher which is selected from the group consisting of cellulose acetate phthalate, hydroxypropylmethyl cellulose phthalate, polyvinyl acetate phthalate, carboxymethylethylcellulose, Eudragit L (poly(methacrylic acid, methylmethacrylate), 1:1 ratio; MW (No. Av. 135,000 - USP Type A) or Eudragit S (poly(methacrylic acid, methylmethacrylate), 1:2 ratio MW (No. Av. 135,000 - USP Type B) and mixtures thereof.

The enteric coating agent may also include an inert processing aid in an amount in the case of the tabletted form from 15 to 55wt% and preferably 20 to 45wt% based on the total weight of the acid resisting component and the inert processing aid. In the case of the pelleted form the inert processing aid is preferably in an amount from 5 to 50wt% and most preferably 10-20wt%. The inert processing aids include finely divided forms of talc, silicon dioxide, magnesium stearate etc. Typical solvents which may be used to apply the acid resisting component-inert processing aid mixture include isopropyl alcohol,

acetone, methylene chloride and the like. Generally the acid resistant component-inert processing aid mixture will be applied from a 5 to 20wt% of acid resisting component-inert processing aid mixture based on the total weight of the solvent and the acid resistant component-inert processing aid.

In the case of both the tabletted form and the pelleted form of the invention omeprazole or a pharmaceutically equivalent salt is used in the core.

10 In the tabletted formulation the omeprazole may comprise from 5 to 70wt% and preferably 10 to 30wt% of the granulation. In the pelleted form the Omeprazole may comprise from 10 to 50wt% and preferably 10 to 20wt% of the drug layer composition.

15 A surface active agent is used in both the tabletted and the pelleted form of the invention. The surface active agent may be any pharmaceutically acceptable, non-toxic surfactant. Suitable surface active agents include sodium lauryl sulfate,

20 polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 80 and the like. The surface active agent may be present at a level of from 0.1 to 5wt%. In the case of the tabletted form the surface active agent is preferably 0.20 to 2.0wt% based on the total weight of the granulation. In the pelleted form the surface

25 active agent is preferably 0.20 to 2.0wt% of the total weight of the drug layer composition.

The binder is used in both the tabletted and the pelleted form of the invention. The binder may be any pharmaceutically acceptable, non-toxic pharmaceutically acceptable binder. The binder is preferably a water soluble polymer of the group consisting of polyvinyl alcohol, polyvinylpyrrolidone, methylcellulose, hydroxypropyl cellulose, hydroxymethyl cellulose and

30 the like. A water soluble binder is preferred which is applied from an aqueous medium such as water at a level

35

of from 0.1 to 10wt% and preferably from 0.25 to 7.5wt% of binder based on the total weight of the granulation.

In the case of the tabletted form of the invention a granulation is formed by contacting the alkaline
5 agent, the omeprazole, the surface active agent and the binder with a medium which may comprise any low viscosity solvent such as water, isopropyl alcohol, acetone, ethanol or the like. When fluids such as water are employed, this will usually require a weight of
10 fluid which is about three times the weight of the dry components of the coating composition.

After the granulation is formed and dried, the granulation is tabletted and the tablets are directly coated with the enteric coating agent. A color
15 imparting agent may be added to the enteric coating agent mixture or a rapidly dissolving seal coat containing color may be coated over the enteric coating agent layer provided that the seal coat is compatible with and does not affect the dissolution of the enteric
20 coating layer. The rapidly dissolving seal coat may comprise Opadry pink which comprises approximately 91wt% hydroxypropyl methylcellulose (E-6), color and 9wt% polyethylene glycol which is applied as a 8-15%w/w solution in purified water. In addition the color may
25 be provided as Chromateric which is available from Crompton & Knowles. This product contains water, talc, TiO_2 , triethyl citrate, propylene glycol, synthetic red iron oxide, potassium sorbate, xanthan gum, sodium citrate and synthetic yellow iron oxide. If desired,
30 conventional sugar based seal coats may be used which contain FDA certified dyes.

In the case of a pelleted form the invention is preferably based on pellets having a core forming inert component which may comprise a starch or sugar sphere
35 such as non-pareil sugar seeds having an average size of 14 to 35 mesh, preferably about 18 to 20 mesh. The

core forming inert component is coated with a formulation which comprises Omeprazole, a surface active agent, a filler, an alkaline material that is either lysine or arginine and a binder, which are collectively referred to as the drug layer composition.

The core forming inert component is employed at 1:1 to 5:1 and preferably from 2:1 to 3:1 weight ratio to the drug layer composition.

The cores are formed by spraying the non-pareil seeds with an aqueous or non-aqueous suspension which contains the alkaline agent, the omeprazole, the surface active agent and the binder. The suspension medium may comprise any low viscosity solvent such as water, isopropyl alcohol, acetone, ethanol or the like.

When fluids such as water are employed, this will usually require a weight of fluid which is about seven times the weight of the dry components of the coating composition.

After the cores are dried, the cores are coated with the enteric coating agent. A color imparting agent may be added to the enteric coating agent mixture or a rapidly dissolving seal coat over the enteric coating agent layer provided that the seal coat is compatible with and does not affect the dissolution of the enteric coating layer.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

Examples 1 to 5 describe a tableted form of the invention and Example 6 describes a pelleted form of the invention.

EXAMPLE 1

Granulation.

A granulation containing omeprazole is formed in a fluid bed coater using a top spray granulation forming

suspension containing omeprazole, micronized to 95% less than 15 microns, 5%w/w of the total amount of L-arginine, polyvinyl pyrrolidone, sodium lauryl sulfate and purified water which is sprayed onto a mixture of microcrystalline cellulose, 95%w/w of the total amount of L-arginine and sodium starch glycolate. The formulation for making the granulation has the following composition:

10	povidone, USP (Plasdone K90)	100.0g
	sodium starch glycolate	100.0g
	sodium lauryl sulfate, NF/USP	6.0g
	microcrystalline cellulose (AvicelPH101)	965.6g
	L-arginine, USP/FCC	1020.0g
15	omeprazole, USP (micronized)	340.0g
	purified water, USP	1100.0g

Tabletting.

The granulation is tabletted into tablets containing 20mg of omeprazole by first mixing the omeprazole granules with glyceryl monostearate:

omeprazole granules	118.0g
glyceryl monostearate (Myvaplex)	6.0g

Tabletting tools: 0.2812"
 target weight : 124mg/tab
 target hardness : 7Kp
 LOD of granules : less than 3%

Enteric coating.

An enteric coating is applied to prepare enteric coated tablets as follows:

35	omeprazole tablets (prepared above)	124.0g
	hydroxypropyl methylcellulose phthalate	

14.7g
talc 4.2g
5 acetyl tributyl citrate 2.9g
acetone 148.0g
isopropyl alcohol 148.0g
10 The solid coating materials were dissolved in the acetone and isopropyl alcohol and this solution was coated onto the omeprazole tablets using a perforated pan
Seal coat:
15 A seal coat was applied to the enteric coated tablets as follows:
Enteric coated tablet 146.0g
Opadry II pink 4.5g
20 Water 450.0g
The seal coat was applied onto the enteric coated omeprazole tablets using a perforated pan coater.

25 EXAMPLE 2

Granulation.

A granulation containing omeprazole is formed in fluid bed coater using a top spray granulation forming
30 suspension containing omeprazole, micronized to 95% less than 15 microns, 2.68%w/w of the total amount of L-arginine, polyvinyl pyrrolidone, polysorbate 80 and purified water which is sprayed onto a mixture of microcrystalline cellulose and 95.0%w/w of the total
35 amount of L-arginine. The formulation for making the granulation has the following composition:

	mg/tablet
povidone, USP (Plasdone K90)	5.88
polysorbate 80 (Tween 80)	0.58

L-arginine, USP/FCC	60.0
omeprazole, USP (micronized)	20.0
microcrystalline cellulose (Avicel PH102)	25.54
purified water, USP	n/a

5

Tabletting.

The granulation is tabletted into tablets containing 20mg of omeprazole by first mixing the omeprazole granules with glyceryl monostearate:

10 omeprazole granules	112.0mg
glyceryl monostearate (Myvaplex)	6.8mg
crospovidone XL	16.2mg

Tabletting tools: 0.2812"

target weight : 135mg/tab

15 target hardness : 7Kp

LOD of granules : less than 3%

Enteric coating.

An enteric coating was applied to prepare enteric coated tablets as follows:

omeprazole tablets (prepared above)	135.0mg
25 Eudragit L30D-55	14.0mg
color (Chromateric)	7.0mg
1M NaOH (to adjust pH to 5.0)qs	na
30 Purified water qs	na

The solid coating materials were dispersed in the water and this mixture was coated onto the omeprazole tablets using a perforated pan.

35

EXAMPLE 3**Granulation.**

A granulation containing omeprazole is formed in

fluid bed coater using a top spray granulation forming suspension containing omeprazole, micronized to 95% less than 15 microns, 5.0%w/w of the total amount of L-arginine, polyvinyl pyrrolidone, sodium lauryl sulfate and purified water which is sprayed onto a mixture of microcrystalline cellulose and 95.0%w/w of the total amount of L-arginine. The formulation for making the granulation has the following composition:

	mg/tablet
10 povidone, USP (Plasdone K90)	5.0
sodium lauryl sulfate	0.3
L-arginine, USP/FCC	60.0
omeprazole, USP (micronized)	10.0g
microcrystalline cellulose (AvicelPH102)	24.7
15 purified water, USP	n/a

Tabletting.

The granulation is tabletted into tablets containing 10mg of omeprazole by first mixing the omeprazole granules with glyceryl monostearate:

omeprazole granules	100.0mg
glyceryl monostearate (Myvaplex)	5.0mg
sodium starch glycolate	5.0mg

Tabletting tools: 0.2812"

25 target weight : 110mg/tab
target hardness : 7Kp
LOD of granules : less than 3%

Enteric coating.

30 The tablets were coated with the same enteric coating that was applied to the tablets in Example 2.

EXAMPLE 4

35 Granulation.

A granulation containing omeprazole is formed in

fluid bed coater using a top spray granulation forming suspension containing omeprazole, micronized to 95% less than 15 microns, 5.0%w/w of the total amount of L-arginine, polyvinyl pyrrolidone, sodium lauryl sulfate and purified water which is sprayed onto a mixture of microcrystalline cellulose and 95.0%w/w of the total amount of L-arginine. The formulation for making the granulation has the following composition:

	mg/tablet
10 povidone, USP (Plasdone K90)	5.88
polysorbate 80	0.60
L-arginine, USP/FCC	60.0
omeprazole, USP (micronized)	20.0
crospovidone XL	5.88
15 microcrystalline cellulose	25.54
purified water, USP	n/a

Tabletting.

The granulation is tabletted into tablets containing 20mg of omeprazole by first mixing the omeprazole granules with glyceryl monostearate:

omeprazole granules	117.9mg
glyceryl monostearate (Myvaplex)	6.1mg

Tabletting tools: 0.2812"

25 target weight : 124mg/tab

target hardness : 7Kp

LOD of granules : less than 3%

Enteric coating.

30 The tablets were coated with the same enteric coating that was applied to the tablets in Example 1.

EXAMPLE 5

35 The granulation of Example 1 was prepared and tabletted into tablets containing 20.0mg of omeprazole.

These tablets were coated as follows:

Enteric coating.

An enteric coating was applied to prepare enteric coated tablets as follows:

5	omeprazole tablets (prepared above)	126.00mg
	Eudragit L30D-55	17.00mg
10	1M NaOH (to adjust pH to 5.0)qs	na
	acetyl tributyl citrate	1.70mg
15	talc	3.80mg
	polysorbate 80	1.50mg
	Purified water qs	na

20

The solid coating materials were dispersed in the water and this mixture was coated onto the omeprazole tablets using a perforated pan. A seal coat was applied using the procedure of Example 1.

25

EXAMPLE 6

In the case of a pharmaceutical formulation with a pelleted omeprazole core, the core is comprised of omeprazole, a surface active agent, a filler, an alkaline material and a binder.

Omeprazole activated pellets (sodium free) are prepared as follows: 13.650 kg of Purified water is dispensed into a suitably sized stainless steel container. L-Arginine Base (0.210 kg), Lactose Anhydrous, NF (1.75 kg) and Povidone (Plasdone ®K-90) (0.056 kg) is added to the purified water while homogenizing at full speed (about 5,000 rpm). Homogenizing is continued until the materials are completely dissolved. Polysorbate 80, NF (0.044 kg) is added to the solution while homogenizing at a lower speed (700 -3300 rpm) to avoid excess foaming.

The material is homogenized until dissolved completely. Half of the solution (7.855 kg) is transferred into a 5-10 gallon stainless steel container. The original container is hereafter
5 referred to as "container A" and the new container is henceforth referred to as "container B." Micronized omeprazole 95% less than 15 microns (0.980 kg) is added to container A while homogenizing at a lower speed (700-3300 rpm) to avoid excess foaming. The Omeprazole
10 is allowed to disperse into the solution completely and then homogenized for another 10 minutes. The homogenizer is replaced with a mechanical stirrer and the suspension is continuously stirred throughout the coating process. When approximately three fourth of
15 the omeprazole suspension in container A is consumed, 0.980 kg of micronized omeprazole is added to container B while homogenizing at a lower speed (700-3300) to avoid excess foaming. The Omeprazole is allowed to disperse in the solution completely and homogenization
20 is continued for another 10 minutes. The homogenizer is replaced with a mechanical stirrer and the suspension is continuously stirred throughout the coating process. 9.98 kg of sugar spheres are added to a fluidized bed coater and preheated until the
25 product reaches 40-45°C. The drug suspension from containers A and B are sprayed onto the spheres. The atomization pressure is between 1.5 to 3.5 bar and the pump rate is 2-100 ml/min. The spray rate does not exceed 20ml/min in the first two hours to avoid
30 agglomeration of the sugar spheres. The coating suspension is transferred to a smaller container to facilitate stirring when the surface of the coating suspension reaches the stirring blade. After the coating suspension has been consumed the pump is
35 stopped and the fluidization is continued in the fluidized bed coater with the heat off until the

product temperature drops below 32°C.

The pellets are then transferred to a fluidized bed coater into a 50°C oven (45-55°C). The pellets are dried until the moisture content of the pellets is not
5 more than 2.5%. The pellets are separated into different size fractions by using a SWECO Separator equipped with 14 and 24 mesh screens. The pellets are collected in doubled polyethylene lined plastic containers and stored with desiccant.

10 ENTERIC COATING PROCESS

10.844 kg of isopropyl alcohol, USP is dispensed into a suitably sized stainless steel container. 10.844 kg of acetone is added to the isopropyl alcohol. 1.683 kg of hydroxypropyl methylcellulose
15 phthalate (Hypromellose 55, Substitution type 200731) and cetyl alcohol, NF (0.084 kg) are added to the solution while homogenizing at full speed until all the materials are dissolved completely. The homogenizer is then removed and replaced with a
20 mechanical stirrer.

Talc (1.683 kg) is added while stirring. The talc is mixed until fully dispersed in the solution and the mixing is continued throughout the entire coating process. A fluidized bed coater is preheated
25 to 32°C. The omeprazole active pellets (11.550 kg) are loaded into the fluidized bed coater and preheated until the temperature reaches 30°C.

The coating suspension is sprayed on the pellets using a product temperature of 25-35°C, an atomization
30 pressure of 1.5 to 3.0 bar and a pump rate of 200-300 ml/min. The coating suspension is transferred to a smaller container to facilitate stirring when the surface of the coating suspension reaches the stirring blade.

35 After the coating suspension has been consumed the coated pellets are dried in a fluidized bed

coater for 20 minutes using the same coating conditions except lowering the atomization pressure to 2 bars or below. The coated pellets are discharged into double polyethylene bags. The pellets are separated into different size fractions by using a SWECO separator equipped with 14 and 24 mesh screens. The pellets which are larger than 14 mesh and smaller than 24 mesh are rejected. The pellets that passed through the 14 mesh and retained on the 24 mesh are retained in polyethylene bags.

BLENDING

Omeprazole enteric coated pellets (Sodium Free), blended are prepared as follows:

14.400 kg of omeprazole enteric coated pellets (Sodium free) are charged into a blender. Talc, USP (0.225 kg) is sprinkled on top of the pellet bed and then blended at 28 rpm for 5 minutes. 0.2 to 0.5 grams of each sample is withdrawn into separate vials from the blender. The blended pellets are unloaded into plastic containers lined with double polyethylene. The excess talc is screened off using a SWECO separator equipped with a 24 mesh screen. The pellets are collected in containers lined with double polyethylene bags and stored with desiccant.

ENCAPSULATION

An encapsulation room is prepared in which the relative humidity is in the range of 35-65% and the temperature is in the range of 15-25°C. Omeprazole enteric coated pellets (Sodium Free) blended are encapsulated using the following equipment and guidelines. A capsule machine model MACOFAR MT-20 is prepared for the procedure placing the machine setting at 4, using capsule machine size part 1, capsule magazine 1. The target filled capsule weight is 457.15 mg. If the total weight is not within 3% of the target weight, further adjustment must be

performed. Capsule fill verification is performed at twenty minutes intervals on ten individual capsules.

Acceptable capsules are collected in containers lined with double polyethylene bags and placed under
5 desiccant.

While certain preferred and alternative embodiments of the invention have been set forth for purposes of disclosing the invention, modifications to the disclosed embodiments may occur to those who are
10 skilled in the art. Accordingly, the appended claims are intended to cover all embodiments of the invention and modifications thereof which do not depart from the spirit and scope of the invention.

15